

## RALEIGH HAND TO SHOULDER CENTER

3701 Wake Forest Rd, Suite 100  
Raleigh, North Carolina 27609  
Telephone: (919) 872-3171  
Fax: (919) 872-6739



### Post-Operative Instructions

1. Please keep your hand elevated above your heart at all times for at least \_\_\_\_\_ days to reduce swelling. Keep a pillow under the hand when sleeping. Prolonged use of an arm sling is not recommended as it holds the hand below the heart.
2. Keep the bandage dry. Cover the bandage with a plastic bag secured with tape or a rubber band while showering.
3. If the fingers are not immobilized by the bandages, you should gently make a fist and straighten the fingers several times per day to decrease stiffness. However, if the fingers are immobilized by a plaster splint, you should rest the hand and avoid movement of the fingers.
4. Call the office if you have severe pain, severe swelling, bleeding, vomiting, fever greater than 101 ° F, wound drainage, or concerns for infection. Some amount of pain, swelling, and bruising is normal after surgery.
5. Call the office if your fingertips become numb or pale. Numbing medicine is often used which causes temporary numbness in the hand for several hours on the day of surgery.
6. If the bandage feels too tight, it may be gently loosened. If this does not help, please notify us. Take care not to move the plaster splint, if one is included in the bandage.
7. If you can take NSAIDs, alternate the pain medication with ibuprofen or naproxen to reduce pain. Wean off the opioid medication as soon as possible. Plan ahead to avoid running out of meds at night and on weekends as these will not be refilled after regular office hours.
8. Do not smoke or chew tobacco. Nicotine can reduce blood flow to the hand and delay healing.
9. Important instructions regarding your post-op bandage:
  - Do not remove the bandage until follow-up appointment.
  - Please remove your bandage \_\_\_\_\_ days after surgery. You may gently clean the area with soap and water, but avoid submerging the incision under water. Then dry the area and cover the incision with dry sterile gauze or a Band-Aid. This should be repeated once daily. Do not apply topical ointments such as Neosporin.
  - Remove the splint while leaving the underlying gauze in place starting the morning after surgery. Move the wrist back and forth several times a day to reduce stiffness. Replace the splint at bedtime. Keep the bandage dry.

Special Instructions: